




# Meniscectomy of the Knee



The meniscus is a commonly injured structure in the knee. The injury can occur in any age group. In younger people the meniscus is fairly tough and rubbery, and tears usually occur as a result of a fairly forceful twisting injury. In older people, the meniscus grows weaker with age, and meniscal tears may occur as a result of a fairly minor injury.

The meniscus is a half moon shaped piece of cartilage that lies between the weight bearing joint surfaces of the thigh and the shin, and is attached to the lining of the knee joint. There are two menisci in a normal knee; the outside one is called the lateral meniscus and the inner one is called the medial meniscus.

The menisci play an important role as a shock absorber in the knee joint, protecting the cartilage that lies on the surface of the bones from impact. The cartilage surface is a tough, very slick material that allows the surfaces to slide against one another without damage to either surface. This ability of the meniscus to spread out the force on the joint surface as we walk is important because it protects the cartilage from excessive forces occurring in anyone area on the joint surface. Without the meniscus, the concentration of force into a small area on the cartilage can damage the surface, leading to degeneration over time. The menisci also cup the joint surfaces of the thigh and therefore provide some degree of stabilization to the knee.

## Meniscal Tears:

There are two different mechanisms for tearing a meniscus.

**Traumatic tears** result from a sudden load being applied to the meniscal tissue that is severe enough to cause the meniscal cartilage to fail and let go. These usually occur from a twisting injury or a blow to the side of the knee that causes the meniscus to be pushed against and compressed.

**Degenerative meniscal tears** are best thought of as a failure of the meniscus over time. The meniscus becomes less elastic and complaint, and as a result may fail with only minimal trauma (such as just getting down into a squat). Sometimes there are no memorable injuries or violent events that can be blamed as the cause of the tear.

## Signs and Symptoms:

The most common problem caused by a torn meniscus is pain. The pain may be felt along the joint line where the meniscus is located or may be more vague and involve the whole knee. Any twisting, squatting or impacting activities will pinch the meniscus tear or flap and cause pain. Often the pain may improve with rest after the initial injury, but as soon as aggressive activity is attempted the pain recurs.

Swelling of the joint may occur although meniscal tears by themselves usually don't cause a large, tensely swollen knee. Typically, low level swelling sets in the next day after the injury and is associated with stiffness and limping.

If the torn portion of the meniscus is large enough, locking may occur. Locking simply refers to the inability to completely straighten out the knee. Locking occurs when the fragment of torn meniscus gets caught in the hinge mechanism of the knee, and will not allow the leg to straighten completely. The torn fragment actually acts like a wedge to prevent the joint surfaces from moving.

There are long term effects of a torn meniscus as well. The constant rubbing of the torn meniscus on the cartilage may cause wear and tear on the surface, leading to degeneration of the joint. The knee may swell with use and become stiff and tight. This is usually because of fluid accumulating inside the knee joint.

## Treatment:

Initial treatment for a torn meniscus usually is directed towards reducing the pain and swelling in the knee. You may be asked to attend physiotherapy to reduce the pain and swelling and improve the range of movement. If the knee is locked and cannot be straightened out, surgery may be recommended as soon as reasonably possible to remove the torn portion that is caught in the knee joint. Once a meniscus is torn, it will most likely not heal on its own. If the symptoms continue, surgery will be required to either remove the torn portion of the meniscus or to repair the tear.

## Surgery:

Meniscal surgery is done using the arthroscope. Small incisions are made in the knee to allow the insertion of a small TV camera into the joint. Through another small incision, special instruments are used to remove or the torn portion of meniscus while the arthroscope is used to see what is happening.

In some cases the meniscus tear can be repaired. Sutures are then placed into the torn meniscus until the tear is repaired. Repair of the meniscus is not possible in all cases. Young people with relatively recent meniscal tears are the most likely candidates for repair. Degenerative type tears in older people are not usually repairable.

# What is involved for you as the patient

- Healthy patients are admitted on the morning of their surgery. You should inform your Surgeon and Anaesthetist of any medical conditions or previous medical treatment as this may affect your operation.
- It is extremely important that there are no cuts, scratches, pimples or ulcers on your lower limb as this greatly increases the risk of infection. Your surgery will be postponed until the skin lesions have healed. You should not to shave or wax your legs for one week prior to surgery. • After the operation you will be required to stay in hospital for the day. Overnight stay may be required due to the affects of the anaesthetic or an inability to manage crutches.
- Physiotherapy is commenced immediately post operatively. A physiotherapist will supervise exercises, walking and weight bearing. You should attend to your physiotherapist once or twice in the first week, but do your exercises daily. Sedentary and office workers may return to work approximately 2-3 days following surgery. Most patients should be walking normally 7 days following surgery although there is considerable patient to patient variation.
- Should the left knee be involved then driving an automatic car is possible as soon as pain allows. Should the right knee be involved driving is permitted when you are able to walk without crutches. You must not drive a motor vehicle whilst taking severe pain killing medications.
- Return to vigorous activities will be determined by the extent of the damage to your meniscus and the amount of meniscus that required removal. If minimal damage was present then you may return to vigorous activities after 6 weeks. If significant damage was present then you may be advised to avoid impact loading activities in order to prevent the onset of early arthritis developing within the joint.